arXiv:1006.3069v1 [math.NT] 15 Jun 2010

Preprint

SOME SUPER CONGRUENCES INVOLVING BINOMIAL COEFFICIENTS

HUI-QIN CAO AND ZHI-WEI SUN*

Abstract. Let $p > 3$ be a prime. We show that

$$T_{p-1} \equiv \left(\frac{p}{3}\right) 3^{p-1} \pmod{p^2},$$

where the central trinomial coefficient T_n is the constant term in the expansion of $(1+x+x^{-1})^n$. We also prove three congruences conjectured by Sun one of which is as follows:

$$\sum_{k=0}^{p-1} \binom{p-1}{k}\binom{2k}{k}((-1)^k - (-3)^{-k}) \equiv \left(\frac{p}{3}\right)(3^{p-1}-1) \pmod{p^3}.$$

1. Introduction

As usual we set $\mathbb{N} = \{0, 1, 2, \ldots\}$ and $\mathbb{Z}^+ = \{1, 2, 3, \ldots\}$.

Let $A, B \in \mathbb{Z}$. The Lucas sequences $u_n = u_n(A, B)$ $(n \in \mathbb{N})$ and $v_n = v_n(A, B)$ $(n \in \mathbb{N})$ are defined by

$$u_0 = 0, u_1 = 1, \text{and } u_{n+1} = Au_n - Bu_{n-1} \ (n = 1, 2, \ldots)$$

and

$$v_0 = 2, v_1 = A, \text{and } v_{n+1} = Av_n - Bv_{n-1} \ (n = 1, 2, \ldots).$$

The characteristic equation $x^2 - Ax + B = 0$ has two roots

$$\alpha = \frac{A + \sqrt{\Delta}}{2} \quad \text{and} \quad \beta = \frac{A - \sqrt{\Delta}}{2},$$

where $\Delta = A^2 - 4B$. By induction, one can easily get the following formulas:

$$(\alpha - \beta)u_n = \alpha^n - \beta^n \quad \text{and} \quad v_n = \alpha^n + \beta^n.$$

It is well-known that

$$u_p \equiv \left(\frac{\Delta}{p}\right) \pmod{p} \quad \text{and} \quad u_{p-(\frac{\Delta}{p})} \equiv 0 \pmod{p}$$

for any odd prime p not dividing B (see, e.g., Sun [1]), where $(-)$ denotes the Legendre symbol.

Key words and phrases. Congruences, binomial coefficients, Lucas sequences, central trinomial coefficients

2010 *Mathematics Subject Classification.* Primary 11B65; Secondary 05A10, 11A07, 11B39, 11S99.

*Corresponding author. Supported by the National Natural Science Foundation (grant 10871087) of China.

Let $p>3$ be a prime and m be an integer not divisible by p. In a series of papers Z. W. Sun [1] [2] [3] obtained the following congruences involving binomial coefficients and Lucas sequences:

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}}{m^k}\equiv\left(\frac{\Delta}{p}\right)+u_{p-(\frac{\Delta}{p})}(m-2,1)\pmod{p^2},\tag{1.1}$$

$$\sum_{k=0}^{p-1}\binom{p-1}{k}\frac{\binom{2k}{k}}{(-m)^k}\equiv\left(\frac{\Delta}{p}\right)(m-4)^{p-1}+\left(1-\frac{m}{2}\right)u_{p-(\frac{\Delta}{p})}(m-2,1)\pmod{p^2},\tag{1.2}$$

and

$$\sum_{k=0}^{(p-1)/2}\frac{\binom{2k}{k}}{m^k}\equiv\left(\frac{\Delta}{p}\right)+\left(\frac{-m}{p}\right)\overline{m}u_{p-(\frac{4-m}{p})}(4,m)\pmod{p^2},\tag{1.3}$$

where $\Delta=m^2-4m$ and

$$\overline{m}=\begin{cases}1 & \text{if } m\equiv 4\pmod{p},\\ 2 & \text{if } \left(\frac{4-m}{p}\right)=1,\\ 2/m & \text{if } \left(\frac{4-m}{p}\right)=-1.\end{cases}$$

In [2] Sun made a conjecture of three congruences.

In this paper we will confirm Sun's conjectured congruences involving binomial coefficients and Lucas sequences and prove some identities. Now we state our main results.

Theorem 1.1. *Let $p>3$ be a prime. Then*

$$T_{p-1}\equiv\left(\frac{p}{3}\right)3^{p-1}\pmod{p^2}\tag{1.4}$$

and

$$\sum_{k=0}^{p-1}\binom{p-1}{k}\binom{2k}{k}((-1)^k-(-3)^{-k})\equiv\left(\frac{p}{3}\right)(3^{p-1}-1)\pmod{p^3},$$

where $T_n(n\in\mathbb{N})$ are the central trinomial coefficients.

Theorem 1.2. *Let p be a prime with $p\equiv\pm1\pmod{12}$. Then*

$$\sum_{k=0}^{p-1}\binom{p-1}{k}\binom{2k}{k}(-1)^k u_k(4,1)\equiv(-1)^{(p-1)/2}u_{p-1}(4,1)\pmod{p^3}.$$

Theorem 1.3. *Let p be a prime with $p\equiv\pm1\pmod{8}$. Then*

$$\sum_{k=0}^{p-1}\binom{p-1}{k}\binom{2k}{k}\frac{u_k(4,2)}{(-2)^k}\equiv(-1)^{(p-1)/2}u_{p-1}(4,2)\pmod{p^3}.$$

Theorem 1.4. *(a) If $6\mid n$, then*

$$\sum_{k=0}^{n}\binom{n}{k}\binom{2k}{k}\frac{\left(\frac{k}{3}\right)}{4^k}=0.\tag{1.5}$$

(b) If $n \equiv 3 \pmod 6$*, then*

$$\sum_{k=0}^{n} \binom{n}{k}\binom{2k}{k}\frac{3[3|k]-1}{4^k} = 0. \tag{1.6}$$

Theorem 1.5. *(a) If* $4 \mid n$*, then*

$$\sum_{k=0}^{n} \binom{n}{k}\binom{2k}{k}\frac{u_k(2,2)}{(-4)^k} = 0. \tag{1.7}$$

(b) If $n \equiv 2 \pmod 4$*, then*

$$\sum_{k=0}^{n} \binom{n}{k}\binom{2k}{k}\frac{v_k(2,2)}{(-4)^k} = 0. \tag{1.8}$$

Theorem 1.6. *If* $3 \mid n$*, then*

$$\sum_{k=0}^{n} \binom{n}{k}\binom{2k}{k}\frac{u_k(3,3)}{(-4)^k} = 0. \tag{1.9}$$

2. Two important Lemmas

Lemma 2.1. *Let* $m \in \mathbb{Z} \setminus \{0\}$*. Suppose that* $A > 0$*,* $A^2 - 4B \neq 0$ *and* $mA + 4B = 0$*. For every* $n \in \mathbb{N}$ *we have*

$$\sum_{k=0}^{n} \binom{n}{k}\binom{2k}{k}\frac{u_k(A,B)}{m^k} = \frac{d^{n/2}(\alpha^n - (-\beta)^n)}{m^n(\alpha-\beta)}\sum_{k=0}^{\lfloor n/2\rfloor}\binom{n}{k}\binom{n-k}{k}d^{-k} \tag{2.1}$$

and

$$\sum_{k=0}^{n} \binom{n}{k}\binom{2k}{k}\frac{v_k(A,B)}{m^k} = \frac{d^{n/2}(\alpha^n + (-\beta)^n)}{m^n}\sum_{k=0}^{\lfloor n/2\rfloor}\binom{n}{k}\binom{n-k}{k}d^{-k}, \tag{2.2}$$

where $d = 4 + 4m/A$*.*

Proof. For a polynomial $P(x)$, we use $[x^n]P(x)$ to denote the coefficient of x^n in $P(x)$. It's easy to see that

$$\begin{aligned}[x^n]((1+\alpha x)^2 + mx)^n &= [x^n]\sum_{k=0}^{n}\binom{n}{k}(1+\alpha x)^{2k}(mx)^{n-k}\\ &= m^n\sum_{k=0}^{n}\binom{n}{k}\binom{2k}{k}\frac{\alpha^k}{m^k}.\end{aligned}$$

Clearly,

$$\begin{aligned}[x^n]((1+\alpha x)^2+mx)^n &= [x^n](\alpha^2x^2+(2\alpha+m)x+1)^n\\ &= [x^n]\sum_{\substack{a,b,c\geq 0\\ a+b+c=n}}\binom{n}{a,b,c}\alpha^{2a}(2\alpha+m)^b x^{2a+b}\\ &= \alpha^n\sum_{\substack{a,b\geq 0\\ 2a+b=n}}\binom{n}{a,b,a}\left(2+\frac{m}{\alpha}\right)^b\\ &= \alpha^n\sum_{k=0}^{\lfloor n/2\rfloor}\binom{n}{k}\binom{n-k}{k}\left(2+\frac{m}{\alpha}\right)^{n-2k}.\end{aligned}$$

So we obtain that

$$m^n\sum_{k=0}^{n}\binom{n}{k}\binom{2k}{k}\frac{\alpha^k}{m^k}=\alpha^n\sum_{k=0}^{\lfloor n/2\rfloor}\binom{n}{k}\binom{n-k}{k}\left(2+\frac{m}{\alpha}\right)^{n-2k}. \tag{2.3}$$

Similarly,

$$m^n\sum_{k=0}^{n}\binom{n}{k}\binom{2k}{k}\frac{\beta^k}{m^k}=\beta^n\sum_{k=0}^{\lfloor n/2\rfloor}\binom{n}{k}\binom{n-k}{k}\left(2+\frac{m}{\beta}\right)^{n-2k}. \tag{2.4}$$

Since $mA+4B=m(\alpha+\beta)+4\alpha\beta=0$, we have $4+\frac{m}{\alpha}+\frac{m}{\beta}=0$ and

$$\begin{aligned}2+\frac{m}{\alpha}&=2+\frac{2m}{A+\sqrt{A^2-4B}}\\ &=2+\frac{2m(\sqrt{A^2+mA}-A)}{mA}\\ &=\frac{2\sqrt{A^2+mA}}{A}=\sqrt{d}.\end{aligned}$$

Note that $(\alpha-\beta)u_k=\alpha^k-\beta^k$ and $v_k=\alpha^k+\beta^k$. Combining (2.3) and (2.4) we get (2.1) and (2.2) immediately. □

For $n\in\mathbb{N}$ we set $H_n=\sum_{k=1}^n 1/k$.

Lemma 2.2. *Let $p>3$ be a prime and $d\in\mathbb{Z}$ with $p\nmid d$. Then*

$$\begin{aligned}&\sum_{k=0}^{(p-1)/2}\binom{p-1}{k}\binom{p-1-k}{k}d^{-k}\\ \equiv&\left(\frac{\Delta}{p}\right)\left(\frac{1-d^{p-1}}{2}+(d-4)^{p-1}\right)-\frac{d}{4}u_{p-(\frac{\Delta}{p})}(d-2,1) \pmod{p^2},\end{aligned}$$

where $\Delta=d(d-4)$.

Proof. Since $\binom{p-1}{k} \equiv (-1)^k(1-pH_k) \pmod{p^2}$ and $\binom{p-1-k}{k} \equiv (-1)^k\binom{2k}{k} \pmod{p}$, we have

$$\begin{aligned}&\sum_{k=0}^{\frac{p-1}{2}}\binom{p-1}{k}\binom{p-1-k}{k}d^{-k}\\ \equiv&\sum_{k=0}^{\frac{p-1}{2}}(-1)^k(1-pH_k)\binom{p-1-k}{k}d^{-k}\\ =&\sum_{k=0}^{\frac{p-1}{2}}\binom{p-1-k}{k}(-d)^{-k}-p\sum_{k=0}^{\frac{p-1}{2}}H_k\binom{p-1-k}{k}(-d)^{-k}\\ \equiv&\sum_{k=0}^{\frac{p-1}{2}}\binom{p-1-k}{k}(-d)^{-k}-p\sum_{k=0}^{\frac{p-1}{2}}H_k\binom{2k}{k}d^{-k} \pmod{p^2}.\end{aligned}$$

It is well known that

$$u_{n+1}(A,B)=\sum_{k=0}^{n/2}\binom{n-k}{k}A^{n-2k}(-B)^k \quad \text{for} \quad n\in\mathbb{N}.$$

So we have

$$\sum_{k=0}^{\frac{p-1}{2}}\binom{p-1-k}{k}(-d)^{-k}=(-d)^{1-p}u_p(-d,d)=d^{1-p}u_p(d,d).$$

By [S09a, Lemma 2.4] we have

$$2u_p(d,d)-\left(\frac{\Delta}{p}\right)d^{p-1}\equiv u_p(d-2,1)+u_{p-(\frac{\Delta}{p})}(d-2,1) \pmod{p^2}. \tag{2.5}$$

On the other hand, in view of [S09d, (2.6)] we have

$$u_p(d-2,1)-\left(\frac{\Delta}{p}\right)\equiv\left(\frac{d}{2}-1\right)u_{p-(\frac{\Delta}{p})}(d-2,1) \pmod{p^2}. \tag{2.6}$$

Combining the above two congruences we immediately get

$$u_p(d,d)\equiv\left(\frac{\Delta}{p}\right)\frac{(d^{p-1}+1)}{2}+\frac{d}{4}u_{p-(\frac{\Delta}{p})}(d-2,1) \pmod{p^2}.$$

Thus

$$\begin{aligned}\sum_{k=0}^{\frac{p-1}{2}}\binom{p-1-k}{k}(-d)^{-k}\equiv&\left(\frac{\Delta}{p}\right)\frac{d^{p-1}+1}{2d^{p-1}}+\frac{d}{4d^{p-1}}u_{p-(\frac{\Delta}{p})}(d-2,1)\\ \equiv&\left(\frac{\Delta}{p}\right)\frac{d^{p-1}+1}{2d^{p-1}}+\frac{d}{4}u_{p-(\frac{\Delta}{p})}(d-2,1) \pmod{p^2}.\end{aligned}$$

Clearly

$$\begin{aligned}
&p\sum_{k=0}^{(p-1)/2}H_k\binom{2k}{k}d^{-k}\\
\equiv&\sum_{k=0}^{(p-1)/2}\left(1-(-1)^k\binom{p-1}{k}\right)\binom{2k}{k}d^{-k}\\
=&\sum_{k=0}^{(p-1)/2}\frac{\binom{2k}{k}}{d^k}-\sum_{k=0}^{(p-1)/2}\binom{p-1}{k}\frac{\binom{2k}{k}}{(-d)^k}\\
=&\sum_{k=0}^{(p-1)/2}\frac{\binom{2k}{k}}{d^k}-\sum_{k=0}^{p-1}\binom{p-1}{k}\frac{\binom{2k}{k}}{(-d)^k}+\sum_{k=\frac{p+1}{2}}^{p-1}\binom{p-1}{k}\frac{\binom{2k}{k}}{(-d)^k}\\
\equiv&\sum_{k=0}^{p-1}\frac{\binom{2k}{k}}{d^k}-\sum_{k=0}^{p-1}\binom{p-1}{k}\frac{\binom{2k}{k}}{(-d)^k}\ (\mathrm{mod}\ p^2).
\end{aligned}$$

Applying (1.1) and (1.2) we obtain that

$$\begin{aligned}
&p\sum_{k=0}^{(p-1)/2}H_k\binom{2k}{k}d^{-k}\\
\equiv&\left(\frac{\Delta}{p}\right)+u_{p-(\frac{\Delta}{p})}(d-2,1)-\left(1-\frac{d}{2}\right)u_{p-(\frac{\Delta}{p})}(d-2,1)-\left(\frac{\Delta}{p}\right)(d-4)^{p-1}\\
=&\left(\frac{\Delta}{p}\right)(1-(d-4)^{p-1})+\frac{d}{2}u_{p-(\frac{\Delta}{p})}(d-2,1).
\end{aligned}$$

Hence

$$\begin{aligned}
&\sum_{k=0}^{\frac{p-1}{2}}\binom{p-1}{k}\binom{p-1-k}{k}d^{-k}\\
\equiv&\left(\frac{\Delta}{p}\right)\left(\frac{1-d^{p-1}}{2d^{p-1}}+(d-4)^{p-1}\right)-\frac{d}{4}u_{p-(\frac{\Delta}{p})}(d-2,1)\\
\equiv&\left(\frac{\Delta}{p}\right)\left(\frac{1-d^{p-1}}{2}+(d-4)^{p-1}\right)-\frac{d}{4}u_{p-(\frac{\Delta}{p})}(d-2,1)\ (\mathrm{mod}\ p^2).
\end{aligned}$$

The proof is now complete. □

Note that $\left(\frac{-3}{p}\right)=\left(\frac{p}{3}\right)$, $u_{p-(\frac{-1}{p})}(0,1)=0$ and

$$u_{p-(\frac{-3}{p})}(-1,1)=u_{p-(\frac{p}{3})}(-1,1)=(-1)^{p-(\frac{p}{3})-1}u_{p-(\frac{p}{3})}(1,1)=\left(\frac{p-(\frac{p}{3})}{3}\right)=0.$$

Applying Lemma 2.2 with $d=1,2,3$, we get corollary as follows:

Corollary 2.1. *Let $p>3$ be a prime. Then*

$$\sum_{k=0}^{\frac{p-1}{2}}\binom{p-1}{k}\binom{p-1-k}{k}3^{-k}\equiv\left(\frac{p}{3}\right)\frac{3-3^{p-1}}{2}\ (\mathrm{mod}\ p^2).\qquad(2.7)$$

$$\sum_{k=0}^{\frac{p-1}{2}}\binom{p-1}{k}\binom{p-1-k}{k}2^{-k}\equiv\left(\frac{-1}{p}\right)\frac{2^{p-1}+1}{2}\pmod{p^2}.\tag{2.8}$$

$$\sum_{k=0}^{\frac{p-1}{2}}\binom{p-1}{k}\binom{p-1-k}{k}\equiv\left(\frac{p}{3}\right)3^{p-1}\pmod{p^2}.\tag{2.9}$$

3. PROOFS OF THEOREMS 1.1-1.6

Proof of Theorem 1.1. The central trinomial coefficients

$$T_k=\sum_{j=0}^{k}\binom{k}{j}\binom{k-j}{j}\qquad k=0,1,2,\ldots.$$

In view of (2.9) we immediately get (1.4).

Clearly $u_{p-1}(4,3)\equiv 0\pmod{p}$. Apply (2.1) with $n=p-1, A=4, B=3, m=-3$ and $d=4+4m/A=1$. By (2.9), we obtain that

$$\begin{aligned}&\sum_{k=0}^{p-1}\binom{p-1}{k}\binom{2k}{k}(-1)^k u_k(4,3)\\ \equiv&\frac{u_{p-1}(4,3)}{(-3)^{p-1}}\sum_{k=0}^{(p-1)/2}\binom{p-1}{k}\binom{p-1-k}{k}\\ \equiv&\frac{1}{3^{p-1}}\left(\frac{p}{3}\right)3^{p-1}u_{p-1}(4,3)\\ =&\left(\frac{p}{3}\right)u_{p-1}(4,3)\pmod{p^3}.\end{aligned}$$

Note that $u_k(4,3)=3^k-1$ for $k\in\mathbb{N}$. We conclude the proof now. □

Proof of Theorem 1.2. If $p\equiv\pm1\pmod{12}$ then

$$3^{(p-1)/2}\equiv\left(\frac{3}{p}\right)=(-1)^{(p-1)/2}\left(\frac{p}{3}\right)=1\pmod{p}.$$

Therefore $u_{p-1}(4,1)\equiv 0\pmod{p}$ and $3^{p-1}\equiv 2\cdot 3^{(p-1)/2}-1\pmod{p^2}$. Apply (2.1) with $n=p-1, A=4, B=1, m=-1, d=4+4m/A=3$. With the help of (2.7), we obtain that

$$\begin{aligned}&\sum_{k=0}^{p-1}\binom{p-1}{k}\binom{2k}{k}(-1)^k u_k(4,1)\\ \equiv&3^{(p-1)/2}u_{p-1}(4,1)\sum_{k=0}^{(p-1)/2}\binom{p-1}{k}\binom{p-1-k}{k}3^{-k}\\ \equiv&3^{(p-1)/2}\left(\frac{p}{3}\right)\frac{3-3^{p-1}}{2}u_{p-1}(4,1)\\ \equiv&(-1)^{(p-1)/2}3^{(p-1)/2}(2-3^{(p-1)/2})u_{p-1}(4,1)\\ \equiv&(-1)^{(p-1)/2}u_{p-1}(4,1)\pmod{p^3}.\end{aligned}$$

□

Proof of Theorem 1.3. If $p \equiv \pm 1 \pmod 8$ then

$$2^{(p-1)/2} \equiv \left(\frac{2}{p}\right) = 1 \pmod p.$$

Therefore $u_{p-1}(4,2) \equiv 0 \pmod p$ and $2^{p-1}+1 \equiv 2 \cdot 2^{(p-1)/2} \pmod{p^2}$. Put $n = p-1, A = 4, B = 2, m = -2, d = 4 + 4m/A = 2$ in (2.1). Applying (2.8), we obtain that

$$\begin{aligned}&\sum_{k=0}^{p-1}\binom{p-1}{k}\binom{2k}{k}(-1)^k u_k(4,2)\\ \equiv&\frac{2^{(p-1)/2}u_{p-1}(4,2)}{(-2)^{p-1}}\sum_{k=0}^{(p-1)/2}\binom{p-1}{k}\binom{p-1-k}{k}2^{-k}\\ \equiv&\frac{1}{2^{(p-1)/2}}\left(\frac{-1}{p}\right)\frac{1+2^{p-1}}{2}u_{p-1}(4,2)\\ \equiv&(-1)^{(p-1)/2}u_{p-1}(4,2) \pmod{p^3}.\end{aligned}$$

□

Proof of Theorem 1.4. Put $A = B = 1, m = -4$. Then $mA + 4B = 0$ and

$$\alpha^2 + \alpha\beta + \beta^2 = (\alpha+\beta)^2 - \alpha\beta = 0.$$

Thus $\alpha^3 = \beta^3$. Consequently $\alpha^n - (-\beta)^n = 0$ provided that $6 \mid n$ and $\alpha^n + (-\beta)^n = 0$ if $n \equiv 3 \pmod 6$. Notice that $u_k(1,1) = (-1)^{k-1}\left(\frac{k}{3}\right)$ and $v_k(1,1) = (-1)^k(3[3|k]-1)$. With the help of Lemma 2.1 we get the equations (1.5) and (1.6) promptly. □

Proof of Theorem 1.5. Set $A = B = 2, m = -4$. It is easy to see that $mA + 4B = 0$ and

$$\alpha^2 + \beta^2 = (\alpha+\beta)^2 - 2\alpha\beta = 0,$$

that is $\alpha^2 = -\beta^2$. So we have $\alpha^n - (-\beta)^n = 0$ provided that $4 \mid n$ and $\alpha^n + (-\beta)^n = 0$ if $n \equiv 2 \pmod 4$. By Lemma 2.1 we get the equations (1.7) and (1.8) promptly. □

Proof of Theorem 1.6. Let $A = B = 3, m = -4$. Clearly $mA + 4B = 0$ and

$$\alpha^2 - \alpha\beta + \beta^2 = (\alpha+\beta)^2 - 3\alpha\beta = 0.$$

Therefore $\alpha^3 = -\beta^3$. Hence $\alpha^n - (-\beta)^n = 0$ provided that $3 \mid n$. Applying (2.1) we obtain the equation (1.9) immediately. □

REFERENCES

[1] Z. W. Sun, *Binomial coefficients, Catalan numbers and Lucas quotients*, preprint, arXiv:0909.5648. `http://arxiv.org/abs/0909.5648`.

[2] Z. W. Sun, *On sums of binomial coefficients modulo p^2*, preprint, arXiv:0910.5667. `http://arxiv.org/abs/0910.5667`.

[3] Z. W. Sun, *Binomial coefficients, Catalan numbers and Lucas quotients(II)*, Sci. China Math. **53** (2010), in press. `http://arxiv.org/abs/0911.3060`.

[4] Z. W. Sun and R. Tauraso, *On some new congruences for binomial coefficients*, Int. J. Number Theory, to appear. `http://arxiv.org/abs/0709.1665`.

[5] Z. W. Sun and R. Tauraso, *New congruences for central binomial coefficients*, Adv. in Appl. Math. **45**(2010), 125–148.

(Hui-Qin Cao) Department of Applied Mathematics, Nanjing Audit University, Nanjing 211815, People's Republic of China
E-mail address: `caohq@nau.edu.cn`

(Zhi-Wei Sun) Department of Mathematics, Nanjing University, Nanjing 210093, People's Republic of China
E-mail address: `zwsun@nju.edu.cn`